

Ear, Nose & Throat (ENT) and Ophthalmology

Tip Sheet for ICD-10



| Diagnosis | Documentation Requirements | | |
|---|---|--|--|
| Cataracts, Age Related | Document type: -Incipient -Cortical -Anterior or posterior Subcapsular polar -Nuclear -Morgagnian or combined forms | | |
| Cholesteatoma | Document laterality: -Right -Left -Bilateral | Specify site: -External ear -Middle ear | Middle ear must be further identified: -Attic -Tympanum -Mastoid -Diffuse |
| Diabetic Eye Disease | Document diabetes type: -Type 1 -Type II -Long-term insulin use for Type II | Document nonproliferative diabetic retinopathy as: -Mild -Moderate -Severe -With or without macula -Edema | Document abnormal blood sugar control as: -Inadequately controlled -Poorly controlled -Out of control |
| Drug Underdosing | Document type: - Intentional versus - Unintentional | Document reason for underdosing, such as: -Financial hardship or - Age related dementia | |
| Dysphagia | Document phase: - Oral - Oropharyngeal - Pharyngeal - Pharyngo-esophageal | Document if sequela of nontraumatic hemorrhage: specify type: - Subarachnoid - Intracerebral - Intracranial | Document if sequela of: - Cerebral infarction - Cerebrovascular disease |
| Eustachian Salpingitis | Document type: - Acute - Chronic | There is no code for Eustachian Tube Dysfunction (ETD) | Specify: Eustachian tube obstruction: -intrinsic or extrinsic cartilaginous or osseous -Patulous Eustachian tube |
| Hearing Loss | Document laterality: - Right - Left - Bilateral | Document type: - Conductive - Sensorineural - Mixed - Noise-induced - Ototoxic (specify agent, e.g. Carboplatin, Erythromycin, etc.) | Document any hearing differences in each ear Document when impacted cerumen is present |
| Hyperthyroidism (Thyrotoxicosis) | Document: - Diffuse goiter - Single or multinodular - Ectopic thyroid tissue - Due to ingestion of excessive thyroid hormone | Document: - With thyrotoxic crisis/storm - Without thyrotoxic crisis/storm. | Document any thyrotoxicosis factitia (drug- induced) |
| Mastoiditis | Document laterality: - Right - Left - Bilateral | Document acuity: - Acute (specify if acute conditions is with OR without complications) - Chronic | Detail if present: - Subperiosteal abscess - Petrositis - Postauricular fistula List any underlying disease (e.g., infectious or parasitic diseases) |

Ear, Nose & Throat (ENT) and Ophthalmology

Tip Sheet for ICD-10



| Diagnosis | Documentation Requirements | | |
|---|---|---|--|
| Neoplasms | Document specific site: - External upper or lower lip - Inner upper or lower lip - Superior, posterior, lateral, or anterior wall of nasopharynx | Differentiate between primary and secondary (metastatic) site | Secondary site: - Document primary site and if it is still present |
| Otitis Externa | Document laterality: - Right - Left - Bilateral Document acuity: - Acute - Chronic | Document infective type: - Abscess - Cellulitis - Malignant - Diffuse - Hemorrhagic - Swimmers ear - Cholesteatoma | Document non-infective type: - Actinic - Chemical - Contact - Eczematoid - Reactive |
| Otitis Media | Document acuity: - Acute - Subacute - Chronic - Recurrent Document laterality: - Right - Left - Bilateral | Document: - Spontaneous rupture of eardrum - Myringitis Document Type: - Serous - Mucoid - Allergic - Atticantral - Tubotympanic | Differentiate: - Non-suppurative - Suppurative Document any tobacco use, abuse, dependence, or exposure Document underlying diseases (viral infection, influenza, etc.) |
| Rhinitis | Document acuity: - Acute - Chronic Document: - With sore throat - Allergic with asthma (specify if exacerbation or status asthmaticus) | Document type: - Allergic (due to food or pollen, seasonal, nonseasonal, perennial) - Infective - Pneumococcal - Syphilitic - Tuberculosis - Vasomotor | Document any tobacco use, abuse, dependence, or exposure |
| Sinusitis | Document acuity: - Acute - Chronic - Acute Recurrent | Document location: - Maxillary - Frontal - Ethmoidal - Sphenoidal - Pansinusitis | Document any tobacco use, abuse, dependence, or exposure |
| Tinnitus, Dysfunction of Eustachian Tube | Document laterality: - Right - Left - Bilateral | Document: - Underlying disease or injury Document: - Circumstances around any injury, medical misadventure or mishap causing an ear condition | Document any tobacco use, abuse, dependence, or exposure Document any hearing differences in each ear |
| Tobacco Use Disorder | Document type: - Cigarettes - Chewing tobacco - Other Delineate between: - Tobacco use/abuse - Tobacco dependence | Document state of dependence: - In remission - With withdrawal - Without withdrawal | Document if used during pregnancy, childbirth, Puerperium Describe history, including product and time |

Ear, Nose & Throat (ENT) and Ophthalmology

Tip Sheet for ICD-10



Diagnosis Documentation Requirements

| | | | |
|--|--|--|---|
| Tonsillitis | Document acuity: - Acute - Chronic (with or without adenoiditis) - Recurrent | Document type/cause: - Diphtheritic - Hypertrophic (with or without adenoiditis) - Staphylococcal - Tuberculosis - Vincent's | Differentiate between: - Tonsillitis - Peritonsillar abscess Document any tobacco use, abuse, dependence, or exposure |
| Tympanic Membrane Disorders | List any associated otitis media Specify any related trauma and associated circumstances surrounding the injury | Specify perforations: - Central - Attic - Multiple - Total - Other | Specify myringitis as: - Acute - Bullous - Chronic Specify atrophic tympanic membrane as flaccid or non-flaccid |
| Upper Respiratory Disorders | Document acuity: - Acute - Chronic - Subacute - Recurrent | Document site: - Maxillary - Pharynx - Larynx - Other | Specify infectious agent, when known (e.g. Streptococcus, influenza A) Document any hypertrophy or obstruction Document underlying or associated conditions (e.g. COPD, abscess, etc.) |
| Procedures | Documentation Requirements | | |
| Adenoidectomy & Tonsillectomy | Document whether: - Partial removal - Complete removal | | |
| Aspiration | Document location, laterality and approach | Document whether drainage device was used | |
| Biopsy | Document location, laterality and approach | | |
| Parotidectomy | Document location and laterality | Document whether partial or complete removal and the approach | |
| Mechanical Ventilation | Document: - Consecutive hours of ventilation (< 24hrs, 24-96 hrs, or > 96hrs) | Document support type: - CPAP - IPAP - IPPB | Differentiate between respiratory assistance (Extracorporeal) or respiratory performance |
| Removal Impacted Cerumen | Document approach: - Percutaneous - Via natural or artificial opening - Endoscopic | | |